Proposal for verifying Wigner's rotation with massive spin-$1/2$ particles at non-relativistic speeds

Veiko Palge*
Graduate School of Information Science, Nagoya University, Nagoya 464-8601, Japan

Jacob Dunningham†
Department of Physics and Astronomy, University of Sussex, Brighton BN1 9QH, United Kingdom

Yuji Hasegawa‡
Atominstitut, Vienna University of Technology, Stadionalle 2, 1020 Vienna, Austria

arXiv:1604.03389v1 [quant-ph] 12 Apr 2016

In quantum physics Wigner's rotation is commonly regarded as confirmed by the Thomas precession in a hydrogen like atom. In this paper we show that a direct experimental verification of Wigner's rotation is in principle accessible in the regime of non-relativistic velocities at $2 \cdot 10^3$ m/s and propose an experiment using thermal neutrons. The experiment can be carried out in a laboratory and it provides a test of relativity in the quantum domain.

Introduction.—Over the last decades a large number of experiments have been carried out to verify the foundations of physics. Their significance lies in providing direct tests (in contrast to mere effects) of particular properties of physical systems to a high degree of accuracy. Wigner rotation first came to the fore in the context of atomic physics when Thomas showed that the hydrogen atom needs a relativistic correction due to a phenomenon that came to be called the Thomas precession [1]. In a nutshell, the Wigner rotation is a phenomenon that if a physical system undergoes non-collinear Lorentz boosts, then it is rotated relative to the original frame. Measurements of atomic spectra provide indirect confirmation of the Wigner rotation.

In this paper we show that a direct experimental verification of Wigner rotation is in principle accessible in the regime of non-relativistic velocities at $2 \cdot 10^3$ m/s in an implementation with thermal neutrons. It is intriguing that one can detect observable effects at speeds which are three orders of magnitude smaller than the speed of electron in the hydrogen atom. From a geometric point of view the experiment can be regarded as explicitly verifying the curvature of the relativistic velocity space—a feature that sets apart the relativistic and non-relativistic theories of spacetime.

Wigner rotation.—Wigner rotation originates in the fact that the subset of pure Lorentz boosts does not form a subgroup of the Lorentz group. The combination of two boosts $\Lambda(v_1)$ and $\Lambda(v_2)$ is in general a boost *and* a rotation,

$$\Lambda(v_2)\Lambda(v_1) = W(\omega)\Lambda(v_3), \tag{1}$$

where $W(\omega) \in \mathrm{SO}(3)$ is the Wigner rotation with angle ω around the axis $\hat{n} = \hat{v}_2 \times \hat{v}_1$ orthogonal to the plane specified by v_1 and v_2. Wigner rotation is a truly relativistic phenomenon both in the conceptual and quantitative senses. The latter means that in a simple scenario consisting of two boosts the rotation angle becomes significant at large velocities. For $v_{1,2} = 0.5c$ when velocities are orthogonal the rotation is $8°$, while at $v_{1,2} = 0.1c$ it is barely measurable at $0.3°$. However, one can achieve a measurable effect at low velocities by noting another property of the Wigner rotation, namely, that it is *cumulative*. From a geometric point of view, in the Wigner representation a massive relativistic spin-$1/2$ particle with mass m is described by $\mathbb{C}^2$-valued square integrable functions on the mass hyperboloid. Equivalently, it is easier to work in the velocity space, which is a hyperbolic space with constant Gaussian curvature $K = 1$. In this picture, a boost corresponds to parallel transportation of the quantum state along a geodesic. Since the velocity space is curved, parallel transportation of spin around a closed loop will in general result in a net rotation. The angle $\Delta\alpha$ between the initial and final vectors is given by [2]

$$\Delta\alpha = \int_U K \, \mathrm{d}S, \tag{2}$$

where U is the region whose boundary ∂U corresponds to the path of the particle.

Experiment.—Consider a setup shown in Fig. 1 where a spin-$1/2$ particle with mass m and momentum $|p\rangle$ comes in from the left with the initial spin prepared in state $|\uparrow\rangle$. We assume that the momentum and spin lie in the plane of the figure. The particle then follows a path around the circle where it undergoes a series of small non-collinear boosts, each of which produces a tiny Wigner rotation. The total Wigner angle $\omega \equiv \Delta\alpha$ accumulated as a result of a single orbital revolution around the circle at constant speed v can be calculated with Eq. (2). Using polar coordinates the volume ele-

* Email: veiko.palge@gmail.com
† Email: J.Dunningham@sussex.ac.uk
‡ Email: hasegawa@ati.ac.at



Figure 1. Experimental setup for measuring Wigner rotation. Path of the particle is shown as solid line. The particle with momentum $|p\rangle$ and spin $|\uparrow\rangle$ enters from the left and orbits k times around the circle. It then exits towards the measurement apparatus M which measures the direction of spin. Each orbital revolution around the circle produces spin rotation by ω, resulting in the total rotation of $k\omega$.

ment of the velocity space can be written as

$$\mathrm{d}S = \frac{u}{(1-u^2)^{3/2}}\,\mathrm{d}u\,\mathrm{d}\phi. \tag{3}$$

For a circle with radius v we get [3]

$$\omega(v) = \int_0^{2\pi}\int_0^{v} \frac{u}{(1-u^2)^{3/2}}\,\mathrm{d}u\,\mathrm{d}\phi = 2\pi\left[\gamma(v)-1\right]. \tag{4}$$

While this angle is still too small for detection in practice, the effect can be amplified by letting the particle travel around the circle k times: this produces the total rotation of $k\omega$. The initial and final states of the particle are related by the evolution

$$|p\rangle\otimes|\uparrow\rangle \longrightarrow |p\rangle\otimes U(k\omega)\,|\uparrow\rangle, \tag{5}$$

where $U(\omega) = \exp\left(-\mathrm{i}\omega\hat{n}\boldsymbol{\sigma}/2\right)$ is a rotation by ω around unit vector $\hat{n}$ orthogonal to the plane of the circle, with $\boldsymbol{\sigma} = (\sigma_x, \sigma_y, \sigma_z)$ the vector of Pauli matrices. A measurement M carried out on the final spin state $U(k\omega)\,|\uparrow\rangle$ shows the difference between the directions of the initial and final spins. Note that we have assumed that the particle undergoes maximal Wigner rotation. This is because we specified above that the initial spin lies in the plane of the circle or, equivalently, in the plane of the boosts around the circle.

Thermal neutrons.—A possible implementation of the experiment at non-relativistic speeds is in the context of neutron polarimetry [4–6]. Let us analyze the magnitude of the effect in this setting. Our system will consist of thermal neutrons traveling at $v = 2\cdot 10^3\,\mathrm{m/s}$ and having a mean lifetime $\tau = 887\,\mathrm{s}$ or about $15\,\mathrm{minutes}$. Suppose the circle has radius r, then the system makes $v/2\pi r$ revolutions in unit time $1\,\mathrm{s}$. We obtain the total rotation ω_T by multiplying with the duration t of the experiment and the angle $\omega(v)$ per one revolution given in Eq. (4),

$$\omega_T(v,r,t) = \frac{v}{2\pi r}\cdot t\cdot\omega(v). \tag{6}$$

Plot of $\omega_T(v,r,t)$ for thermal neutrons with $t=\tau$ as well as longer times is shown in Fig. 2. We see that a total



Figure 2. Dependence of the total Wigner rotation ω_T on the radius r of circle for a particle with $v = 2\cdot 10^3\,\mathrm{m/s}$, and duration of experiment t is τ (red), 2τ (blue), 3τ (green) and 4τ (black).

spin rotation of about $1.1°$ will be observed for mean lifetimes when the guiding ring has radius $r = 2\,\mathrm{mm}$. This is close to the precision of $1°$ achievable in a standard configuration [7]. However, since the total rotation is proportional to the duration of experiment, the effect will be twice as large at about $2.2°$ for experiments that last twice as long, and even larger for longer times. Longer experiments need to take into account loss of signal due to loss of neutrons through beta decay and although this is a restriction it is not a critical issue.

Conclusion.—We have discussed an experiment that allows to verify Wigner's rotation in the non-relativistic regime. The speeds involved are significantly lower than that of an electron in a hydrogen (like) atom which scales as $1/137\,c$. Another advantage lies in that the scheme provides a direct test by measuring spin rotation. This is in contrast to atomic spectroscopy, which is commonly regarded as a confirmation of Wigner's rotation in the quantum domain and where one observes an effect of Wigner's rotation—the spectrum which arises due to a contribution from the Wigner rotation. Although mean lifetime leads to a spin rotation of $1.1°$, the parameters determining the outcome are tunable and enable larger effects as well. An experiment that lasts about an hour yields a larger angle of $4.4°$. In the same vein, a ring with smaller radius gives more spin rotation since radius is inversely proportional to the total rotation ω_T in Eq. (6).

Acknowledgments.—This work was financially supported by the Japan Society for the Promotion of Science (JSPS) Grant-in-Aid for JSPS Fellows No. 25-03757.

[1] L. H. Thomas, Nature **117**, 514 (1926).
[2] T. Frankel, *The Geometry of Physics: An Introduction* (Cambridge University Press, 2012).
[3] This coincides with the result obtained in [8] using a slightly different method.
[4] Y. Hasegawa and G. Badurek, Physical Review A **59**, 4614 (1999).
[5] S. Sponar, J. Klepp, C. Zeiner, G. Badurek, and Y. Hasegawa, Physics Letters A **374**, 431 (2010).
[6] B. Demirel, S. Sponar, and Y. Hasegawa, New Journal of Physics **17**, 023065 (2015).
[7] Better precision has been obtained in a particular experiment [9].
[8] J. A. Rhodes and M. D. Semon, American Journal of Physics **72**, 943 (2004).
[9] Y. Hasegawa, C. Schmitzer, H. Bartosik, J. Klepp, S. Sponar, K. Durstberger-Rennhofer, and G. Badurek, New Journal of Physics **14**, 023039 (2012).